

TT Computed tomography of the body

The computed tomography (CT) device performs cross-section or 'slice' imaging, in which the image is formed with the help of X-rays. For this reason, the study is not suitable for pregnant women.

How to prepare for the procedure?

Before the procedure, you can eat and take medication normally. You must drink 1 litre of water 1-2 hours before the scan. The toilet should not be used for approximately 45 minutes prior to the examination, to allow urine to fill the bladder during the examination.

If your doctor thinks it is necessary to check your kidney function before the procedure, they will order a blood test. Metformin medication does not need to be stopped before the procedure.

If you are, or think you may be, pregnant or if you are allergic to X-ray contrast agents, please tell this to the referring doctor and please contact the X-ray department before the day of the examination.

If you have previously had a mild reaction to an Iodine contrast agent, a contrast examination can often be done with pre-administered cortisone medication. If you have previously had a severe contrast reaction, the study will be performed without contrast.

How is the procedure done?

During the procedure it is important for you to keep absolutely still and follow the breathing instructions given to you by the Radiographer. The CT table moves during the scan and the Radiographer will be able to see and hear you throughout the scan.

An intra-venous contrast agent may be used during the CT scan. In general, contrast agents do not have side effects. However, if you notice anything unusual about your health after the examination, please contact the examination department or on-call doctor.

It is a good idea to set aside about an hour for the procedure. Urgent on-call examinations are also carried out from the ward, so your start time may be delayed.

What about after the procedure?

The procedure does not require follow-up at the hospital. After the procedure it is recommended to drink plenty of water. The referring unit is responsible for reporting the results of the procedure.