

Fecal culture 3 (F-BaktVi3, 3584)

The test is done to identify the bacteria that cause toxin-mediated food poisoning, *Bacillus cereus*, *Clostridium perfringens* or *Staphylococcus aureus*. The test is performed only to study the etiology of food poisoning epidemic. Contact the laboratory, tel. 02-313 1675 before sending the epidemic sample

Materials

1. Plastic or cardboard container
2. Copan Eswab 490CE.A swab and tube (Fig. 1)
3. Plastic bag and sticker for name



Fig. 1 Copan eSwab

Procedure

1. Defecate into the container or on toilet paper.
2. Roll the swab with the cotton end in the feces, especially in parts of the feces containing blood or mucus. Avoid taking too much of the feces for the sample.
3. If the feces sample is liquid, hold the cotton end of the swab in the sample for a moment.
4. Transfer immediately after sampling the swab into the transport tube that contains some liquid. Note! Do not pour the liquid out from the tube. Break the swab at the indicated breaking point and close the stopper.
5. **Write your name, social security number, sampling date and sampling time on the sticker that came with the materials from the lab.**
6. **Attach the sticker onto the transportation tube.**
7. Place the transportation tube into the plastic bag.
8. Keep the Copan eSwab tube at room temperature

Take the Copan eSwab sample tube to the laboratory on the day after the sampling, at the latest. If the transportation is delayed the sample must be kept at refrigerator temperature

Please contact the department or outpatient clinic if you have questions regarding the sampling procedure for this examination.

You may ask for the result of this test at the hospital unit where you are being treated.

Reviewed 6.9.2022 MG